

Necessary and Sufficient Condition that the Limit of Stieltjes Transforms is a Stieltjes Transform

Jeffrey S. Geronimo¹ and Theodore P. Hill²

School of Mathematics
Georgia Institute of Technology
Atlanta, GA 30332-0160

Abstract

The pointwise limit S of a sequence of Stieltjes transforms (S_n) of real Borel probability measures (P_n) is itself the Stieltjes transform of a Borel p.m. P if and only if $iyS(iy) \rightarrow -1$ as $y \rightarrow \infty$, in which case P_n converges to P in distribution. Applications are given to several problems in mathematical physics.

Key words and phrases: real Borel probability measure, convergence in distribution, Stieltjes transform, Lévy continuity theorem, Akhiezer-Krein theorem, weak convergence of probability measures.

Lévy's classical continuity theorem says that if the pointwise limit of the characteristic functions of a sequence of real Borel probability measures (P_n) exists, then the limit function φ is itself the characteristic function for a probability measure P if and only if φ is continuous at zero, in which case $P_n \rightarrow P$ in distribution. The purpose of this note is to prove a direct analog of Lévy's theorem for Stieltjes transforms, complementing those for other representing functions in [HS] and [HK], and to give several examples of applications.

Throughout this note, \mathbb{R} and \mathbb{C} denote the real and complex numbers, respectively; p.m. and s.p.m. denote Borel probability measures, and sub-probability (mass ≤ 1) measures, respectively, on \mathbb{R} ; and s.p.m.'s (μ_n) *converge vaguely* to a s.p.m. μ [C, p. 80], if there exists a dense subset D of \mathbb{R} such that for all $a, b \in D$, $a < b$, $\mu_n((a, b]) \rightarrow \mu((a, b])$. (Thus if $(\mu_n), \mu$ are p.m.'s, vague convergence is equivalent to convergence in distribution.)

¹ Research partially supported by NSF Grant DMS-9970613.

² Research partially supported by NSF Grant DMS-9971146 and Göttingen Academy of Sciences Gauss Professorship (Fall 2000).

Definition 1. The *Stieltjes transform* S_P of a p.m. P is the function $S_P : \{\text{Im}(z) > 0\} \rightarrow \mathbb{C}$ given by

$$S_P(z) = \int_{-\infty}^{\infty} \frac{1}{w - z} dP(w).$$

A basic property of Stieltjes transforms, which has important applications in the theory of moments (cf. [A], [S1], [ST]) and in mathematical physics (), is that they are a representing class for finite measures.

Lemma 1. For s.p.m.'s P and Q , $P = Q$ iff $S_P = S_Q$.

Proof. Follows immediately from the Stieltjes transform inversion formula [A, p. 125]. \square

Just as limits of characteristic functions of p.m.'s are in general not characteristic functions, and limits of Hardy-Littlewood functions or expected-extrema functions are not in general Hardy-Littlewood or expected-extrema functions [HK], limits of Stieltjes transforms are not always Stieltjes transforms, as the next easy example shows.

Example 1. For $n = 1, 2, \dots$, let $P_n = \delta_{(n)}$, the Dirac point mass at n . Then $S_{P_n}(z) = (n - z)^{-1}$ for all n and all z with $\text{Im}(z) > 0$, so $\lim_{n \rightarrow \infty} S_{P_n}(z) \equiv 0$, which is clearly not the Stieltjes transform for any p.m. P (see Lemma 2 below).

On the other hand, just as with Lévy's theorem, the limit of Stieltjes transforms is itself a Stieltjes transform if and only if it satisfies one single universal limit condition. The next theorem is the main result of this note.

Theorem 1. Suppose that (P_n) are real Borel probability measures with Stieltjes transforms (S_n) , respectively. If $\lim_{n \rightarrow \infty} S_n(z) = S(z)$ for all z with $\text{Im}(z) > 0$, then there exists a Borel probability measure P with Stieltjes transform $S_P = S$ if and only if

$$\lim_{y \rightarrow \infty} iyS(iy) = -1, \tag{1}$$

in which case $P_n \rightarrow P$ in distribution.

Corollary 1. If P , (P_n) are real Borel p.m.'s with Stieltjes transforms S , (S_n) , respectively, then $P_n \rightarrow P$ in distribution if and only if $S_n \rightarrow S$ pointwise.

Proof of Corollary. If $S_n \rightarrow S$, then $P_n \rightarrow P$ in distribution by Theorem 1. Conversely, suppose that $P_n \rightarrow P$ in distribution. Since $f_z(w) := (w - z)^{-1}$ is continuous and bounded in w for fixed z in $\{\text{Im}(z) > 0\}$, then $\text{Im}(f_z)$ and $\text{Re}(f_z)$ are also continuous and bounded, so by the basic equivalence of convergence in distribution of p.m.'s and convergence of integrals of bounded continuous functions [C, Theorem 4.4.2], $\int \text{Im}(f_z) dP_n \rightarrow \int \text{Im}(f_z) dP$ and $\int \text{Re}(f_z) dP_n \rightarrow \int \text{Re}(f_z) dP$, so $S_n(z) \rightarrow S(z)$. \square

To facilitate the proof of Theorem 1, two additional lemmas are useful, which are stated here for ease of reference.

Lemma 2. *Let $S : \{\text{Im}(z) > 0\} \rightarrow \mathbb{C}$ be analytic. Then there exists a p.m. P with $S_P(z) = S(z)$ for all z with $\text{Im}(z) > 0$ if and only if (1) holds and*

$$\text{Im}(S(z)) > 0 \quad \text{for all } z \text{ with } \text{Im}(z) > 0. \quad (2)$$

Proof. By the classical Akhiezer-Krein theorem [A, p. 93], $S = S_P$ for some finite Borel measure P if and only if: S is analytic in $\{\text{Im}(z) > 0\}$; S satisfies (2); and

$$\sup_{y \geq 1} |yS(iy)| < \infty. \quad (3)$$

Suppose P is a p.m. with $S = S_P$. The Akhiezer-Krein theorem implies that (2) holds, and (1) follows immediately from the definition of S_P . Conversely, suppose that S is analytic and satisfies (1) and (2). Since $yS(iy)$ is continuous in y , (1) easily implies (3), so by the Akhiezer-Krein theorem again, there is a finite Borel measure P with $S_P = S$. Since clearly $\lim_{y \rightarrow \infty} [-iyS_P(iy)] = \text{mass}(P)$, (1) implies that P is a p.m. \square

Lemma 3. *Let \mathcal{F} be a family of functions analytic in a connected open domain D . If for each compact $K \subset D$ there exists a constant $M(K) < \infty$ such that*

$$(4) \quad |f(z)| \leq M(K) \quad \text{for all } z \in K \text{ and all } f \in \mathcal{F},$$

then all pointwise limits of functions in \mathcal{F} are also analytic in D .

Proof. ([H, Theorem 15.2.3]). \square

Proof of Theorem 1. If $\lim S_n = S = S_P$ for some p.m. P , then (1) follows by Lemma 2.

Conversely, suppose that $S = \lim_n S_n$ satisfies (1). Let $\mathcal{F} = \{\bigcup S_n\}$, and for $K \subset D := \{\text{Im}(z) > 0\}$, let $d(K) = \inf\{\|y - z\| : y \in \mathbb{R}, z \in K\}$, the smallest distance from K to the real line. Clearly $0 < d(K) < \infty$ for all compact $K \subset D$, and $M(K) = 1/d(K)$ satisfies (4), so Lemma 3 implies that $S = \lim S_n$ is analytic in D . By Lemma 2, $\text{Im}(S_n(z)) > 0$ for all $z \in D$, so $\text{Im}(S(z)) \geq 0$ for all $z \in D$. Suppose, by way of contradiction to (2), that $\text{Im}(S(z_0)) = 0$ for some $z_0 \in D$. Since S is analytic, $\text{Im}(S)$ and $\text{Re}(S)$ are harmonic on D [K, p. 590]. By the maximum principle [K, p. 760], a non-constant function which is harmonic in a simply connected bounded open set G has neither a maximum nor a minimum in G , so since $\text{Im}(S(z)) \geq 0$ on G for every simply connected open bounded set G with $z_0 \in G \subset D$, it follows (taking $G_t = \{z \in D : \|z\| < t\}$, and letting $t \rightarrow \infty$) that $\text{Im}(S(z)) \equiv 0$ for all $z \in D$, which contradicts (1). Thus (2) holds, and since S is analytic and (1) holds by assumption, Lemma 2 implies there exists a real Borel p.m. P with $S_P = S$.

For the convergence in distribution conclusion, suppose that $S_n = S_{P_n} \rightarrow S_P$ pointwise in D for p.m.'s (P_n) , P . By the Helly selection theorem [C, Theorem 4.3.3], there exists a s.p.m. Q and a subsequence (P_{n_k}) of (P_n) such that $P_{n_k} \rightarrow Q$ vaguely. Fix z in D , and let $f_z : \mathbb{R} \rightarrow \mathbb{C}$ be given by $f_z(w) = (w - z)^{-1}$. Since f_z is continuous in w and vanishes at infinity, $\text{Re}(f_z)$ and $\text{Im}(f_z)$ are continuous and vanish at infinity, so it follows by the equivalence of vague convergence of s.p.m.'s and convergence of integrals of continuous functions which vanish at infinity [C, Theorem 4.4.1] that $S_{P_{n_k}}(z) \rightarrow S_Q(z)$ as $k \rightarrow \infty$ for all $z \in D$. By hypothesis, $S_{P_n} \rightarrow S_P$, so $S_P = S_Q$, which by Lemma 1 implies that $P = Q$. Since every vaguely convergent subsequence of (P_n) thus converges to P , this implies [C, Theorem 4.3.4] that P_n converges vaguely to P , that is, since (P_n) and P are p.m.'s, P_n converges to P in distribution. \square

Sketch of Alternative Proof. (B. Simon [S2]). The functions $\{S_{P_n}\}$ are Herglotz functions, so the limit S is Herglotz, and pointwise convergence $S_n \rightarrow S$ implies weak convergence for the measures $(1+x^2)^{-1}dP_n$ to a measure P on $[-\infty, \infty]$, where $(1+x^2)^{-1}dP$ is finite. Given that $S(iy) \rightarrow 0$ as $y \rightarrow \infty$, $P(\{-\infty, \infty\}) = 0$, so it follows using the fact

that $S_{P_n} \rightarrow S_P$, the Herglotz representation theorem, and the monotone convergence theorem, that P is a p.m. Then weak convergence of P_n to P can be shown given the weak convergence of the measures when multiplied by $(1 + x^2)^{-1}$. (For related ideas, see pp. 129–130 in [S1]). \square

References

- [A] Akhiezer, N. (1965). *The Classical Moment Problem*, Hafner, New York.
- [AK] Akhiezer, N. and Krein, M. (1962). *Some Questions in the Theory of Moments, Translation of Mathematical Monographs*, vol. 2, Amer. Math. Society, Providence.
- [C] Chung, K. (1968). *A Course in Probability*, 2nd Ed., Academic Press, San Diego.
- [HK] Hill, T. and Krenzel, U. (2001). “Levy-like continuity theorems for convergence in distribution,” accepted for publication in *Proceedings of the Göttingen Academy of Sciences*.
- [HS] Hill, T. and Spruill, M. (1994). “On the relationship between convergence in distribution and convergence of expected extremes,” *Proc. Amer. Math. Soc.* **121**, 1235–1243.
- [H] Hille, E. (1959). *Analytic Function Theory*, vol. II, Chelsea, NY.
- [K] Kreyszig, E. (1962). *Advanced Engineering Mathematics*, Wiley, New York.
- [S1] Simon, B. (1998). “The classical moment problem as a self-adjoint finite difference operator,” *Adv. in Math.* **137**, 82–203.
- [S2] Simon, B. (2001). Private communication.
- [ST] Shohat, J. and Tamarkin, J. (1943). “The problem of moments,” *Amer. Math. Soc. Surveys* **1**.