

## A NONSTANDARD DELTA FUNCTION

TODOR TODOROV

(Communicated by James E. West)

**ABSTRACT.** We prove that the Dirac delta distribution has a kernel in the class of the pointwise nonstandard functions.

The purpose of this note is to prove the existence of a nonstandard function  $\Delta: {}^*\mathbb{R}^n \rightarrow {}^*\mathbb{C}$  such that

$$\int_{\mathbb{R}^n} \Delta(x) {}^*\varphi(x) dx = \varphi(0)$$

for all  $\varphi \in C^0$ . Here  $C^0 \equiv C^0(\mathbb{R}^n)$  is the class of the continuous complex-valued functions defined by  $\mathbb{R}^n$ ,  ${}^*\mathbb{R}$  and  ${}^*\mathbb{C}$  are the sets of the nonstandard real and nonstandard complex numbers, respectively, and  ${}^*\varphi: {}^*\mathbb{R}^n \rightarrow {}^*\mathbb{C}$  is the nonstandard extension of  $\varphi$ . For examples of nonstandard functions  $\Delta$  for which (1) holds merely "up to infinitesimals," we refer the reader to one of the many texts on nonstandard analysis, e.g. [2, p. 300]. Recall that there does not exist a standard function  $\Delta$  with the property mentioned above.

In what follows, we shall work in a nonstandard model with a set of individuals  $S$  that contains the complex numbers  $\mathbb{C}$  and degree of saturation  $k$  larger than  $2^\kappa$  for  $\kappa = \text{card } C^0$ . In particular, any polysaturated model of  $\mathbb{C}$  will do [2].

**Notation.** For any  $\varphi \in C^0$ , we define the functional  $F_\varphi: \mathcal{D} \rightarrow \mathbb{C}$  by

$$(2) \quad F_\varphi(f) = \int_{\mathbb{R}^n} f(x)\varphi(x) dx, \quad f \in \mathcal{D},$$

where  $\mathcal{D} \equiv C_0^\infty(\mathbb{R}^n)$  is the class of all  $C^\infty$ -functions on  $\mathbb{R}^n$  with compact support. We write  $\ker F_\varphi$  for the kernel of  $F_\varphi$ . For the nonstandard extension  ${}^*F_\varphi: {}^*\mathcal{D} \rightarrow {}^*\mathbb{C}$  of  $F_\varphi$  for  $\varphi \in C^0$ , we have the \*-integral representation

$$(3) \quad {}^*F_\varphi(f) = \int_{{}^*\mathbb{R}^n} f(x) {}^*\varphi(x) dx, \quad f \in {}^*\mathcal{D}.$$

---

Received by the editors July 24, 1989.

1980 *Mathematics Subject Classification* (1985 Revision). Primary 03H05, 30G10, 46F05, 46F99.

*Key words and phrases.* Nonstandard analysis, nonstandard extension, saturation principle, delta function.

**Lemma.** For any  $k \in \mathbb{N}$  and any  $\varphi_i \in C^0$ ,  $i = 1, 2, \dots, k$ , the system of equations

$$(4) \quad F_{\varphi_i}(f) = \varphi_i(0), \quad i = 1, 2, \dots, k,$$

has a solution  $f$  in  $\mathcal{D}$ .

*Proof.* Consider first the case  $k = 1$  of one equation:

$$(5) \quad F_{\varphi}(f) = \varphi(0).$$

If  $\varphi = 0$ , then any  $f$  in  $\mathcal{D}$  is a solution of (5). If  $\varphi \neq 0$ , then the set  $\Phi \equiv \mathcal{D} - \ker F_{\varphi}$  is nonempty and the function

$$f = \frac{\varphi(0)}{F_{\varphi}(g)} g$$

satisfies (5) for any choice of  $g \in \Phi$ . Assume, now, that the statement is true for  $k-1$ . If  $\varphi_1, \dots, \varphi_k$  are linearly dependent in  $C^0$ , then (4) is equivalent to a system of  $k-1$  equations and, by assumption, has a solution. If  $\varphi_1, \dots, \varphi_k$  are linearly independent, then the sets

$$\Phi_i \equiv \left( \bigcap_{j=1}^k \ker F_{\varphi_j} \right) - \ker F_{\varphi_i}, \quad i = 1, 2, \dots, k,$$

are nonempty [1, vol. 3, Lemma 10, p. 421], and we can pick  $g_i \in \Phi_i$ . Now, the function

$$f = \sum_{i=1}^k \frac{\varphi_i(0)}{F_{\varphi_i}(g_i)} g_i$$

is obviously a solution of (4). The proof is complete.  $\square$

**Proposition.** There exists a nonstandard function  $\Delta \in {}^* \mathcal{D}$  for which (1) holds for all  $\varphi \in C^0$ .

*Proof.* Define the family  $\mathcal{A}_{\varphi}$ ,  $\varphi \in C^0$ , of subsets of  $\mathcal{D}$  by

$$\mathcal{A}_{\varphi} = \{f \in \mathcal{D} : F_{\varphi}(f) = \varphi(0)\},$$

and observe that, by the above lemma, it has the finite intersection property. Hence, by the saturation principle [2, 7.4.2(b), p. 181], the intersection

$$\mathcal{A} \equiv \bigcap_{\varphi \in C^0} {}^* \mathcal{A}_{\varphi}$$

is nonempty, where  ${}^* \mathcal{A}_{\varphi} = \{f \in {}^* \mathcal{D} : {}^* F_{\varphi}(f) = \varphi(0)\}$ . Hence every  $\Delta \in \mathcal{A}$  has the desired property. The proof is complete.  $\square$

#### REFERENCES

1. N. Dunford and J. T. Schwartz, *Linear operators, Part I: general theory*, Interscience Publishers, New York, 1958.
2. K. D. Stroyan and W. A. J. Luxemburg, *Introduction to the theory of infinitesimals*, Academic Press, New York, 1976.