

Intensity interferometry with anyons

Thomas D. Gutierrez*

Department of Physics, University of California, Davis, California 95616

(Received 9 December 2003; published 22 June 2004)

A pairwise correlation function in relative momentum space is discussed as a tool to characterize the properties of an incoherent source of noninteracting Abelian anyons. This is analogous to the Hanbury-Brown Twiss effect for particles with fractional statistics in two dimensions. In particular, using a flux tube model for anyons, the effects of source shape and quantum statistics on a two-particle correlation function are examined. Such a tool may prove useful in the context of quantum computing and other experimental applications where studying anyon sources are of interest.

DOI: 10.1103/PhysRevA.69.063614

PACS number(s): 03.75.Dg, 05.30.Pr, 03.67.Mn, 25.75.Gz

Anyons are particles that exhibit fractional quantum statistics in two dimensions. Existing somewhere between bosons and fermions, anyons have been applied theoretically to a variety of problems such as the fractional quantum Hall effect (FQHE) [1,2], high temperature superconductivity [3], supersymmetry [4,5], and fault-tolerant quantum computing [6]. Experimentally, the quasiparticle excitations seen in the FQHE have been shown to possess anyon properties [7,8].

Fractional spin in two dimensions directly addresses the topological interpretation of quantum statistics. For identical particles in three or more dimensions, the appropriate homotopy group is the permutation group [9] and there are only two homotopy classes. These correspond directly to fermions and bosons.

However, in two dimensions, the corresponding homotopy group is the braid group [9] and an infinite number of homotopy classes are possible. As particles are exchanged in relative position, each successive winding of the particles around each other cannot be smoothly deformed into a finite number of configurations as is the case in higher (>2) dimensions. Each winding corresponds to a separate homotopy class which in turn can be categorized as a different classification of quantum statistics. The term *anyon* was coined by Wilczek [10,11] to describe objects having such topological properties, a physical example of which is a charged particle bound to a magnetic flux tube in two dimensions.

This relationship between topology and statistics was glimpsed by Aharonov and Bohm [12] and later realized, formalized, and expanded by others [10,11,13–16]. Because the homotopy group is the braid group, this makes many multiparticle (>2) anyon problems nearly intractable. An essential analytical difficulty arises because, unlike the case with the permutation group, the multiparticle wave functions for anyons cannot in general be written in a simple way in terms of the single-particle wave functions. Multiparticle anyon wave functions are, in effect, permanently entangled. However, it is exactly this topological property of anyons that makes them robust against decoherence and thus desirable candidates for qubits in fault-tolerant quantum computing [6].

Because quantum interference effects can be sensitive to quantum statistics, it is natural to inquire about the role of anyons in this context. As mentioned, the Aharonov-Bohm effect was an early probe into this fascinating problem. More recently, first order interference effects in a Mach-Zender-style interferometer for non-Abelian anyons in the presence of Aharonov-Bohm flux sources have been derived from applications of the braid group [17].

In this work, intensity interferometry in momentum space, a second-order interference effect, is explored for incoherently emitted, non-interacting Abelian anyons. This is conceptually related to the q -Bose gas interferometry approaches for pions in heavy ion collisions [18,19]. However, q -bosons are not restricted to two dimensions. In that case, the alternative quantum statistics are achieved by directly modifying the commutation relations of the creation and annihilation operators.

Intensity interferometry is a useful tool for characterizing the source geometry of incoherently emitted particles. In addition, the interference effect, expressed in terms of a second-order correlation function, might be regarded as an entanglement measure. Such a tool may prove useful in the context of quantum computing and other experimental applications where studying anyon sources are of interest.

Intensity interferometry was originally developed by Hanbury-Brown and Twiss (HBT) as an alternative to Michelson interferometry to measure the angular sizes of stars in radio astronomy [20]. By correlating intensities rather than amplitudes, the measurement is insensitive to high frequency fluctuations that would normally make Michelson interferometry prohibitive. When applied to classical waves, the HBT effect is essentially a beat phenomenon.

The field of modern quantum optics was spawned when intensity interferometry was quantum mechanically applied to photons rather than classical waves. Now, two- and multiphoton effects are routinely studied in photonics. Two-fermion HBT in two-dimensional condensed matter systems has also been reported [21]. The HBT effect was independently applied to pions and other particles in elementary particle physics in momentum space and is often called the Goldhaber-Goldhaber-Lee-Pais effect in that context [22–24].

Quantum mechanically, the two-particle correlation function is a measure of the degree of independence between a

*Electronic address: tgutierr@lifshitz.ucdavis.edu; URL: <http://nuclear.ucdavis.edu/~tgutierr>

joint measurement of two particles (1 and 2 later). Formally, the correlation function can be written

$$C_2 = \frac{\text{Tr}[\rho a^\dagger a^\dagger a a]}{\text{Tr}[\rho a^\dagger a] \text{Tr}[\rho a^\dagger a]} \sim \frac{P(1,2)}{P(1)P(2)}, \quad (1)$$

where ρ is the density matrix and a^\dagger and a are the creation and annihilation operators for the quanta associated with the appropriate fields of interest. When normalized to the single particle distributions as shown, C_2 is proportional to the relative probability for a joint two-particle measurement as compared to two single-particle measurements. If the measurements are independent, then $C_2=1$. If the measurements are correlated, C_2 deviates from unity. In principle, C_2 can be expressed as a joint measurement as a function of any degree of freedom of interest. In this treatment, we will focus on momentum space correlations.

The second-order correlation function in momentum space is a powerful tool to probe several important properties of a system. C_2 is sensitive to the quantum statistics of the particles as expressed in the commutation relations for the field operators in Eq. (1). Also, as determined by the form of the density matrix, C_2 contains information about the space-time distribution of the particle source in phase space as well as the pairwise interaction and the quantum field configuration.

Noninteracting identical particles can exhibit strong correlations, no correlations, or even anticorrelations depending on the specific field configuration. Possible field configurations for identical bosons include thermal states, coherent states, Fock states, and squeezed states [25]. For example, an incoherent thermal field configuration gives $C_2=2$ for spinless bosons of equal momentum. In contrast, a coherent source of bosons, such as laser light well above the lasing threshold, gives a constant $C_2=1$ for *all* relative momenta.

Geometric information about the source is contained in the shape of the correlation function as a function of the pair's relative momentum. The correlation function typically approaches unity for large momentum differences. For incoherent sources, the relationship between source width and correlation width is essentially limited by the Heisenberg uncertainty principle such that $\delta q \sim 1/R$ where δq is the width of C_2 in relative momentum space and $1/R$ is some characteristic width of the incoherent source generating the pairs.

Keeping in mind the earlier discussion, this paper examines the behavior of C_2 in momentum space for a nonexpanding, nonrelativistic, incoherent source of anyons. An approximation to Eq. (1) is used to highlight some features of intensity interferometry in momentum space. The quantum statistics are "tuned," and, given some simple source functions, the corresponding correlation function is extracted.

In momentum space for an incoherent source, Eq. (1) reduces to the Koonin-Pratt equation [26–28]:

$$C_2(\mathbf{q}) - 1 = \int d\mathbf{r} K(\mathbf{q}, \mathbf{r}) S(\mathbf{r}), \quad (2)$$

where the integration kernel is given by

$$K(\mathbf{q}, \mathbf{r}) = |\Phi_{\mathbf{q}}(\mathbf{r})|^2 - 1. \quad (3)$$

The function $\Phi_{\mathbf{q}}(\mathbf{r})$ is the two-particle wave function in the center of mass frame of the pair, where $\mathbf{q}=(\mathbf{p}_1-\mathbf{p}_2)/2$ is the relative momentum and $\mathbf{r}=\mathbf{r}_1-\mathbf{r}_2$ is the relative separation in that frame. The center of mass motion of the pair is not considered in this treatment. The source function, $S(\mathbf{r})$, is the normalized probability distribution of emitting a particle pair with relative separation \mathbf{r} . The integral is over the entire relative separation space. In this context, incoherent means that the particles are emitted from the source randomly and independently. Moreover, any potential time dependence of the source function in Eq. (2) has been integrated out as in Ref. [28]. This integration eliminates the ability to recover emission-time ordering information between the pairs and places the focus on spatial information only.

Using a flux tube model for anyons, a dynamical approach is used to obtain two-particle nonrelativistic wave functions for anyons in two dimensions [10–12,16]. In this model, the quantum statistics are enforced through an interaction, the strength of which enforces the effective quantum statistics. In this picture, each anyon is analogous to a charged particle bound to a tightly bundled magnetic field in two dimensions. The analogy is not strictly one-to-one in this treatment because the scalar Coulomb potential between the charged pairs is not considered; indeed, the system need not be electromagnetic. Nevertheless, the analogy paints a helpful physical picture.

The tightly bundled magnetic field associated with each anyon, $\mathbf{B}=\nabla \times \mathbf{A}$, points perpendicular to the plane of motion. We can select a gauge such that the vector potential, \mathbf{A} , has the form $A_r=0$ and $A_\phi=\xi/2\pi r$, where ξ is the magnetic flux through the plane. The particles are only permitted in regions of space where the magnetic field of the other particle is zero, so the particles exert no force on each other. Quantum mechanically, the interaction term takes the form of a minimal coupling such that the momentum operator takes the form $\hat{p} \rightarrow (-i\nabla - e\mathbf{A})$. While under the conditions described earlier, this "interaction" exerts no force, it does adjust the phase of the wave functions, permitting interference effects. The relative wave function of the pair can then be used in Eq. (2), in combination with a normalized source distribution, to obtain C_2 .

In the center-of-mass frame, using the Hamiltonian

$$\hat{H} = \frac{\hat{p}_r^2}{m} + \frac{(\hat{p}_\phi - \alpha)^2}{mr^2}, \quad (4)$$

the wave function is obtained for the relative motion of two free anyons in two-dimensional polar coordinates. In Eq. (4), m is the mass of one particle ($m_1=m_2=m$) and α represents the anyon parameter with $0 \leq \alpha \leq 1$, where $\alpha=0(1)$ corresponds to bosons (fermions). In this standard treatment, all of the details of the vector potential, magnetic flux, and gauge choice are included in the tunable parameter α . In Eq. (4), the reduced mass of the identical pair, $\mu=m_1 m_2 / (m_1 + m_2) = m/2$, has already been substituted. In the two-dimensional relative coordinate system of the identical pairs, the x axis has been chosen along the direction of \mathbf{q} .

Applying Eq. (4) to the time-independent Schrödinger equation, $\hat{H}\psi = E\psi$, gives

$$\left[\left(\frac{\partial^2}{\partial r^2} + \frac{1}{r} \frac{\partial}{\partial r} \right) - \frac{1}{r^2} \left(i \frac{\partial}{\partial \phi} + \alpha \right)^2 + q^2 \right] \psi = 0 \quad (5)$$

where $q^2 = mE$ is the square of the relative momentum of the pair (i.e., the energy is that of a free relative particle) and ϕ is the angle between \mathbf{q} and \mathbf{r} . Equation (5) admits solutions of the form

$$\psi(r, \phi) = NJ_{|l-\alpha|}(qr) e^{il\phi}, \quad (6)$$

where N is a normalization constant. We want solutions that correspond to the appropriate symmetric (antisymmetric) boson (fermion) free particle wave function in relative coordinates in the limit of $\alpha=0(1)$. Specifically

$$\Phi_{\alpha=0,1}(r, \phi) = \frac{1}{\sqrt{2}} (e^{i\mathbf{q}\cdot\mathbf{r}} \pm e^{-i\mathbf{q}\cdot\mathbf{r}}) \quad (7)$$

where the $\alpha=0$ case corresponds to the symmetric solution (+) while the $\alpha=1$ case is the antisymmetric solution (-).

Keeping in mind the identity

$$e^{i\mathbf{q}\cdot\mathbf{r}} = e^{qr \cos \phi} = \sum_{n=-\infty}^{+\infty} i^n J_n(qr) e^{in\phi} \quad (8)$$

one can construct partial wave superpositions of Eq. (6) that give Eq. (7) for the limiting cases of $\alpha=0,1$. Namely

$$\Phi_{q,\alpha}(r, \phi) = \sqrt{2} \sum_{l=-\infty(\text{even})}^{+\infty} (i)^{|l-\alpha|} J_{|l-\alpha|}(qr) e^{il\phi}, \quad (9)$$

where l is the orbital angular momentum quantum number. The sum is over partial waves of Bessel functions of the first kind of order $|l-\alpha|$ where l is necessarily an integer, but α is not. The subscript q has been added to the wave function at this stage to remind the reader that we will be integrating over r and ϕ using Eqs. (2) and (3) to obtain a function of q .

Treating the initial pair wave function in Eq. (5) as that of spinless bosons (the ‘‘boson basis’’) simplifies the problem of obtaining the appropriate exchange symmetry for the final two-particle wave function shown in Eq. (9). In constructing Eq. (9), the partial wave expansion using Eq. (6) is symmetrized under the exchange $\phi \rightarrow (\phi - \pi)$. Tuning the anyon parameter will result in a wave function that possesses the correct exchange symmetry for identical indistinguishable pairs. For the fermion and boson cases, Eq. (9), with the help of Eq. (8), reduces to Eq. (7). But for $0 < \alpha < 1$, Eq. (9) is the nontrivial two-anyon free particle wave function.

Equation (2) is used to obtain an expression of C_2 for anyons. While intensity interferometry can be used to image three-dimensional sources, for simplicity we will only consider angle-averaged sources that are a function of r rather than \mathbf{r} . The integration kernel in Eq. (3) is angle averaged in two dimensions and then denoted $K_{0,\alpha}(q, r)$ such that

$$K_{0,\alpha}(q, r) = \left[\frac{1}{2\pi} \int_0^{2\pi} d\phi |\Phi_{q,\alpha}(r, \phi)|^2 \right] - 1. \quad (10)$$

Inserting the expression for the relative wave function from Eq. (9) into Eq. (10) and performing the angle averaging in two dimensions gives

$$K_{0,\alpha}(q, r) = \left[2 \sum_{l=-\infty(\text{even})}^{l=+\infty} |J_{|l-\alpha|}(qr)|^2 \right] - 1. \quad (11)$$

In the limiting case when $\alpha=(0,1)$, Eq. (11) simplifies to, with the help of Bessel function properties

$$K_{0,\alpha=0,1}(q, r) = (-1)^\alpha J_0(2qr), \quad (12)$$

which is, by construction, the same angle-averaged kernels one gets by using Eq. (7) directly. For the more general case when $0 < \alpha < 1$, the sum in Eq. (11) is evaluated numerically to an appropriately converging order.

Two sample source functions are used to illustrate the effects of source shape and size on C_2 :

$$S_g(r) = \frac{e^{-r^2/4r_0^2}}{4\pi r_0^2}, \quad (13)$$

$$S_s(r) = \frac{\Theta(r_0 - r)}{\pi r_0^2}. \quad (14)$$

They represent incoherent pair emission distributions normalized in two dimensions

$$2\pi \int_0^\infty r dr S(r) = 1, \quad (15)$$

and where the effective source width is given by r_0 . The forms of the functions were chosen as examples to represent typical localized incoherent sources.

Integrating Eq. (2) over the azimuthal angle in two dimensions using Eq. (11), and choosing either Eq. (13) or (14) as the source function, gives an angle-averaged expression for C_2 parameterized by α :

$$C_{2,\alpha}(q) - 1 = 2\pi \int_0^\infty r dr K_{0,\alpha}(q, r) S(r). \quad (16)$$

Figure 1 shows $C_{2,\alpha}$ for an incoherent Gaussian source, Eq. (13). Various values of α are displayed. The figures are shown plotted versus the quantity qr_0 , which accounts for the shifting momentum scale upon changing the source size. For example, as the width of the source decreases, the correlation function widens and vice versa.

For the boson and fermion cases, the standard HBT results for incoherent sources are recovered. The top curve in Fig. 1 represents the spin zero case and approaches a value of $C_{2,\alpha=0} = 2$ as $q \rightarrow 0$. Here, it is more likely to measure a pair of noninteracting identical bosons in a state of zero relative momentum than to independently measure each particle with their respective momentum. That is, in the joint measurement, the bosons are not independent. This effect is a function of q and is the result of the Bose-Einstein statistics and the symmetric nature of the wave function. Similarly, the lowest curve represents noninteracting indistinguishable fermions and displays an anticorrelation in the joint measure-




FIG. 1. $C_{2\alpha}$ vs qr_0 from a normalized Gaussian source given by Eq. (13) in two dimensions. The curves represent different values of the anyon parameter α . The top curve is the boson case ($\alpha=0$) and the bottom curve is the fermion case ($\alpha=1$). From top to bottom, $\alpha=0, 0.1, 0.2, 0.3, 0.4, 0.5, 0.6, 0.7, 0.8, 0.9$, and 1. The source function, plotted in arbitrary units, is shown inset vs r/r_0 .

ment at low relative momentum, where $C_{2\alpha=1}=0$ as $q \rightarrow 0$. Similar to the boson case, this is also only due to the quantum statistics and indicates again that a measurement of one particle affects the other quantum mechanically. As the relative momentum increases, the value of $C_{2\alpha}$ approaches unity, indicating that the joint measurement becomes less correlated.

Starting from the top boson curve and moving downward are the correlation curves for various values of the anyon parameter. Like fermions, the anyons display an exclusion principle and always anticorrelate in the limit as $q \rightarrow 0$. However, other than the exclusion at $q=0$, there is a natural trend of decreasing correlation as one interpolates between bosons and fermions for values of $q > 0$. The correlation function also approaches unity for large values of q .

An example of the sensitivity of $C_{2\alpha}$ to the source shape is shown in Fig. 2. The box profile with a sharp edge introduces a ringing structure into the correlation function.

Experimentally, the measured correlation function provides information about the size and shape of the emission




FIG. 2. $C_{2\alpha}$ from a normalized step function source given by Eq. (14). The anyon parameterization is the same as in Fig. 1. The source function, plotted in arbitrary units, is inset plotted vs r/r_0 .

source. Measured correlation functions can be inverted to determine actual source distributions using methods like those discussed in Ref. [28]. Also, information about anyon pairwise interactions can be extracted by comparing measured correlation functions against predictions made using hypothesized interactions in the Hamiltonian. Moreover, the degree of chaoticity of the source, testing the independent emission assumption, can be extracted by looking at the behavior of the correlation at low relative momentum. Finally, the correlation function itself may provide information about the degree of entanglement between pairs versus their relative momentum, a potentially useful tool in the field of quantum computing.

In summary, intensity interferometry in momentum space provides a tool that can be used to experimentally study the properties of anyons and their sources in two dimensions. Further theoretical studies using this framework can be pursued exploring various properties of multianyon systems including anyon interactions, non-Abelian anyon correlations, and higher-order multianyon correlations.

The author would like to thank D. A. Brown, J. L. Klay, S. R. Klein, J. Randrup, R. Vogt, and Z. Yusof for valuable input and stimulating discussions.

-
- [1] B. Halperin, Phys. Rev. Lett. **52**, 1583 (1984).
 [2] J. S. D. Arovas and F. Wilczek, Phys. Rev. Lett. **53**, 722 (1984).
 [3] R. Laughlin, Phys. Rev. Lett. **60**, 2677 (1988).
 [4] M. S. Plyushchay, Mod. Phys. Lett. A **12**, 1153 (1997).
 [5] M. Rausch de Traubenberg and M. J. Slupinski, Mod. Phys. Lett. A **12**, 3051 (1997).
 [6] A. Y. Kitaev, Ann. Phys. (N.Y.) **303**, 2 (2003).
 [7] R. Clark *et al.*, Phys. Rev. Lett. **60**, 1747 (1988).
 [8] J. Simmons *et al.*, Phys. Rev. Lett. **63**, 1731 (1989).
 [9] Y.-S. Wu, Phys. Rev. Lett. **52**, 2103 (1984).
 [10] F. Wilczek, Phys. Rev. Lett. **48**, 1144 (1982).
 [11] F. Wilczek, Phys. Rev. Lett. **49**, 957 (1982).
 [12] Y. Aharonov and D. Bohm, Phys. Rev. **115**, 485 (1959).
 [13] R. M. G. A. Goldin and D. Sharp, J. Math. Phys. **21**, 650 (1980).
 [14] R. M. G. A. Goldin and D. Sharp, J. Math. Phys. **22**, 1664 (1981).
 [15] S. Forte, Rev. Mod. Phys. **64**, 193 (1992).
 [16] A. Khare, *Fractional Statistics and Quantum Theory* (World Scientific, Singapore, 1997).
 [17] B. J. Overbosch and F. A. Bais, Phys. Rev. A **64**, 062107 (2001).
 [18] D. V. Anchishkin, A. M. Gavriliuk, and N. Z. Iorgov, Mod. Phys. Lett. A **15**, 1637 (2000).
 [19] D. V. Anchishkin, A. M. Gavriliuk, and N. Z. Iorgov, Eur. Phys. J. A **7**, 229 (2000).
 [20] R. Hanbury Brown and R. Q. Twiss, Nature (London) **178**, 1046 (1956).
 [21] M. Henny *et al.*, Science **284**, 296 (1999).

- [22] G. Goldhaber, S. Goldhaber, W.-Y. Lee, and A. Pais, *Phys. Rev.* **120**, 300 (1960).
- [23] D. H. Boal, C. K. Gelbke, and B. K. Jennings, *Rev. Mod. Phys.* **62**, 553 (1990).
- [24] G. Baym, *Acta Phys. Pol. B* **29**, 1839 (1998), nucl-th/9804026.
- [25] M. O. Scully and M. S. Zubairy, *Quantum Optics* (Cambridge University Press, Cambridge, 1997).
- [26] S. E. Koonin, *Phys. Lett.* **70B**, 43 (1977).
- [27] S. Pratt, T. Csoergoe, and J. Zimanyi, *Phys. Rev. C* **42**, 2646 (1990).
- [28] D. A. Brown and P. Danielewicz, *Phys. Lett. B* **398**, 252 (1997).