Occam's Razor & The Nature of Scientific Theories

WILLIAM D. STANSFIELD

The early 14th century philosopher William of Occam (Sahakian 1968, p. 116) is credited by historians for proposing that, in interpreting scientific data, the simplest explanation consistent with all the known facts is preferred as a working hypothesis over more complicated ones. In other words, scientific explanations should be pared down (as if using a knife to cut away unnecessary "fat" from a carcass) to the simplest hypothesis as a preliminary orientation for further study of any scientific problem. This guiding principle of scientific methodology has come to be known as "Occam's razor." Genetics is a scientific discipline in which Occam's razor (and other facets of scientific methodology) can be easily explained and grasped by most middle- and high-school students. The following genetics problems can be used to illustrate this point.

In nature, mice usually have a nondescript "mousy" color pattern called "agouti" (named after a South American rodent with the same pattern). The agouti pattern involves a subapical (near the tip) band of yellow pigment in the hair. If the rest of the hair is black, the phenotype is known as "wild-type" (the nor-

mal phenotype found in natural populations). Thus, the wild-type mouse is actually a black-agouti phenotype. If the yellow band appears on an otherwise brown hair (a mutant or rare character in nature), the phenotype is called brown-agouti or cinnamon. Two other mutant types are nonagouti patterns (no yellow band): solid brown (chocolate) and solid black. Students should have, as background information, an understanding of dominant and recessive alleles, multiple alleles, independent assortment, and the statistical inferences that are normally drawn from 3:1 and 9:3:3:1 phenotypic ratios.

Five sets of hypothetical matings will be presented. The parental generation (P) is assumed to consist of genetically pure (homozygous) breeding individuals. Large numbers of progeny are produced in the F₁ and F₂ generations. The F₂ generation is produced by allowing the F₁ population to mate at random. As the data from each set of matings is displayed (e.g., on an overhead projector), the instructor can ask the class to propose suitable explanations. The fifth set of matings should not be revealed until the first four sets have been analyzed.

Set #1

P: wild-type x cinnamon

F₁: all wild-type

F₂: 3/4 wild-type: 1/4 cinnamon

WILLIAM D. STANSFIELD is an Emeritus Professor in the Biological Sciences Department at California Polytechnic State University, San Luis Obispo, CA 93407. He can be reached via e-mail: wstansfi@calpoly.edu.

Inference

A single pair of alleles is segregating in the production of the F_2 generation, with a dominant gene (say D) coding for wild-type and its recessive allele (say d^1) for cinnamon; i.e., $D > d^1$.

Set #2

P: wild-type x black

F_1 : all wild-type

F_2 : 3/4 wild-type: 1/4 black

Inference

Another pair of alleles of the same gene is segregating in the production of the F_2 generation, with dominant gene (D) for wild type, as proposed previously, and a different recessive allele (say d^2) for black. Here we are using Occam's razor for the first time. We are trying to explain both sets of data (Sets #1 and #2) by proposing that three alleles exist at the D-locus in this population of mice. The dominance hierarchy is proposed as follows: $D > (d^1 \text{ and } d^2)$. We do not know, at this point, the relationship (Allchin 2000) among the alleles in parentheses.

Set #3

P: cinnamon x brown

F_1 : all cinnamon

F_2 : 3/4 cinnamon: 1/4 brown

Inference

Another pair of alleles is segregating in the production of the F_2 generation, with the cinnamon gene (d^1) dominant to the recessive brown allele (say d^3). A proposed dominance hierarchy is $D > d^1 > d^3$. At this point, we infer that allele d^2 is lower in the dominance hierarchy than allele D , but its exact relationship to alleles d^1 and d^3 has not been established.

Set #4

P: black x brown

F_1 : all black

F_2 : 3/4 black : 1/4 brown

Inference

The black allele (d^2) is dominant to the brown allele (d^3). A proposed dominance hierarchy is: $D > (d^1 \text{ and } d^2) > d^3$. Now, the only unknown relationship is that between the alleles for cinnamon and black.

Set #5

P: cinnamon x black

F_1 : all wild-type

F_2 : 9/16 wild-type: 3/16 cinnamon: 3/16 black: 1/16 brown

The results of Set #5 will "knock the socks" off of some students. Even the phenotype of the F_1 is obviously inconsistent with the original hypothesis of multiple alleles at a single genetic locus. The most dominant phenotype (wild-type) cannot be produced by matings involving only alleles lower in the dominance hierarchy. Can a more complex hypothesis accommodate all the data? What inference would normally be drawn from a 9:3:3:1 F_2 phenotypic ratio? Answer: Let us assume that the F_1 mice in Set #5 were dihybrids (heterozygous at two independently assorting loci, as on different pairs of chromosomes). We could then define a pair of alleles (A and a) at one locus that governs the agouti (yellow band) and nonagouti (solid color) patterns, respectively; and at an independently assorting locus, another pair of alleles (B and b) that

A+ Performance and Price!

The **PriorLab** laboratory microscope provides a high quality yet economical choice for a broad range of laboratory and classroom needs.

- Inward facing objectives allow easier access to specimens
- Koehler type illumination with field diaphragm & centerable substage
- 6V20W halogen illumination system meets CE requirements
- Newly designed PF optics - providing sharp, flatfield images
- Large built-in graduated mechanical stage
- CCTV and photomicrography capability

Visit us on the web at www.prior.com

PRIOR
Scientific

Prior Scientific Inc.
80 Reservoir Park Drive
Rockland, MA 02370
800-877-2234
•Phone 781-878-8442 •Fax 781-878-8736

governs black and brown, respectively. Set #5 can then be genetically defined as follows:

P: $AAbb$ (cinnamon) $\times$ $aaBB$ (black)

F₁: all $AaBb$ (wild-type)

F₂: 9/16 $A-B-$ (wild-type) : 3/16 $A-bb$ (cinnamon):
3/16 $aaB-$ (black) : 1/16 $aabb$ (brown).

(Note: Each dash in the above genotypes indicates an unknown and irrelevant allele to the production of the phenotype.)

Similarly, the other four sets of matings can be explained in like fashion; it would be good experience for students to do this on their own for homework or perhaps work on it as teams in the classroom. The instructor can take this opportunity to point out that the procedure we have used is precisely the way that scientific knowledge grows. Using Occam's razor (proposing a single genetic locus rather than two loci) as our initial hypothesis was good scientific methodology and worked well in explaining the data through the first four sets of matings. But then came the fifth set of data that (according to Thomas Henry Huxley) exemplifies "The great tragedy of science – the slaying of a beautiful hypothesis by an ugly fact" (Mackay 1977). So, it was back to square one (perhaps a Punnett

square?) in attempts to formulate a more complicated hypothesis that could explain all of the data. All scientific knowledge is tentative in this sense. We may never have all of the data that we want or need to formulate a robust hypothesis. Progress is not made by waiting for more data. Ofttimes we must act on the data and hypotheses that we have, realizing that either may be faulty. Any good scientific hypothesis, however, should point the way to new experiments that will test its falsifiability (Popper 1980, p. 22). A hypothesis that has "stood the test of time" and resisted all attempts at falsification will gradually become accepted as a valid scientific theory.

References

- Allchin, D. (2000). Mending mendelism. *The American Biology Teacher*, 62(9), 633-639.
- Mackay, A.L. (1977). *Scientific Quotations – The Harvest of a Quiet Eye*. New York: Crane, Russak.
- Popper, K. (1980). Science: Conjectures and refutations. In E.D. Klemke, R. Hollinger & A.D. Kline (Eds.), *Introductory Readings in the Philosophy of Science* (pp.19-34). Buffalo, NY: Prometheus Books.
- Sahakian, W.S. (1968). *History of Philosophy*. New York: Barnes & Noble, Inc. College Outline Series.

ASCB to Publish New Journal

Cell Biology Education (CBE), the new education journal owned and published by the American Society for Cell Biology, is accepting manuscript submissions for potential publication in *CBE*'s first issue. *CBE* is an online, quarterly journal publishing peer-reviewed articles on science education. The Journal's initial focus will be on undergraduate education related to cell biology.

For more information about *CBE*, Instructions to Authors, and links to other online resources, see www.cellbioed.org.

Papers can be submitted to cbe@ascb.org or to:

Managing Editor
Cell Biology Education
8120 Woodmont Avenue, Suite 750
Bethesda, MD 20814-2762
Phone: 301-347-9300

Authors choosing to submit manuscripts via email are encouraged to read *CBE*'s Instructions to Authors prior to submission.