

# WIL-POWER™

## SELECTIVE HERBICIDE



**FOR SELECTIVE BROADLEAF WEED CONTROL IN ORNAMENTAL  
LAWNS AND TURF GRASSES, ALSO ON SOD FARMS**

**CONTAINS MCPA, TRICLOPYR AND DICHLORPROP-p**

**CONTROLS:** Dandelion, Chickweed, Black medic, Knotweed, Plantain, Oxalis, Clover, Cocklebur, Thistle  
and many other species of broadleaf weeds; some of which are listed on this label.  
**GET THE OPTICAL ADVANTAGE®**

**ACTIVE INGREDIENTS:**

Dimethylamine Salt of 2-Methyl-4-Chlorophenoxyacetic Acid\* ..... 46.87%  
Triethylamine Salt of 3,5,6-Trichloro-2-Pyridinyloxyacetic Acid\*\* ..... 10.68%  
Dimethylamine Salt of (+)-R-2-(2,4-Dichlorophenoxy)  
propionic Acid\*\*\*† ..... 9.12%

**OTHER INGREDIENTS:** ..... 33.33%  
TOTAL ..... 100.00%

By Isomer specific AOAC Method Equivalent to:

\*2-Methyl-4-Chlorophenoxyacetic Acid ..... 38.27%, 3.72 lbs./gal.  
\*\*3,5,6-Trichloro-2-Pyridinyloxyacetic Acid ..... 7.65%, 0.75 lbs./gal.  
\*\*\*(+)-R-2-(2,4-Dichlorophenoxy) propionic Acid ..... 7.65%, 0.75 lbs./gal.

†CONTAINS THE SINGLE ISOMER FORM OF DICHLORPROP-p

**For Use By Professional Turf Maintenance Personnel,  
Landscaping or Commercial Applicators Only**

EPA REG. NO. 228-411-2935

EPA EST. NO. 228-IL-1

**KEEP OUT OF REACH OF CHILDREN  
DANGER - PELIGRO**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

**SEE INSIDE BOOKLET FOR FIRST AID AND ADDITIONAL  
PRECAUTIONARY STATEMENTS.**

For Chemical Spill, Leak, Fire or Exposure,  
Call CHEMTREC (800) 424-9300

For Medical Emergencies Only, Call (877) 325-1840

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS  
DANGER - PELIGRO**

Corrosive. Causes irreversible eye damage. Harmful if swallowed or absorbed through skin. Do not get in eyes or clothing. Avoid contact with skin. Wash thoroughly with soap and water after handling.

**PERSONAL PROTECTIVE EQUIPMENT (PPE)**

**NON-WPS TURF USE:** Applicators and other handlers who handle this pesticide for any use NOT covered by the Worker Protection Standard (40 CFR Part 170) - in general only agricultural plant uses are covered by the WPS - must wear eye protection (goggles, face shield, or safety glasses) when using this product. It is recommended that safety glasses include front, brow and temple protection.

**WPS SOD FARM USES**

Applicators and other handlers who handle this pesticide for any use covered by the Worker Protection Standard (40 CFR Part 170) - in general, agricultural plant uses are covered - must wear: Long-sleeved shirt and long pants, chemical-resistant gloves made of any waterproof material such as polyethylene or polyvinyl chloride, shoes plus socks and protective eyewear. Some of the materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for Category A on an EPA Chemical-resistant Category Selection Chart. Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining Personal Protective Equipment (PPE). If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

**Engineering Controls Statements:** If the mechanical system is used in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

**USER SAFETY RECOMMENDATIONS**

**USERS SHOULD:**

- Wash hands, face and arms with soap and water before eating, drinking, chewing gum, using tobacco or using the toilet.
- Wash the outside of gloves before removing.

**ENVIRONMENTAL HAZARDS**

Drift or runoff may adversely affect nontarget plants. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. When cleaning equipment, do not pour washwater on the ground; spray or drain over a large area away from wells and other water sources.

Do not contaminate water when disposing of equipment washwaters. Do not apply this product through any type of irrigation system. Do not contaminate water used for irrigation or domestic purposes.

Most cases of ground water contamination involving phenoxy herbicides such as MCPA and Dichlorprop-p have been associated with mixing/loading and disposal sites. Caution should be exercised when handling these phenoxy pesticides at such sites to prevent contamination of groundwater supplies. Use of closed systems for mixing and transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

**FIRST AID**

If in Eyes	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15 to 20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
If Swallowed	<ul style="list-style-type: none"> <li>• Call poison control center or doctor immediately for treatment advice.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• Do not induce vomiting unless told to do so by the poison control center or doctor.</li> <li>• Do not give anything by mouth to an unconscious person.</li> </ul>
If on Skin or Clothing	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15 to 20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>

**HOT LINE NUMBER**

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-877-325-1840 for emergency medical treatment information.

**NOTE TO PHYSICIAN**

Probable mucosal damage may contraindicate the use of gastric lavage.

**NET CONTENTS: 2.5 GALLONS**

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

#### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about Personal Protective Equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 48 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: Coveralls, chemical-resistant gloves made of any waterproof material, shoes plus socks, and protective eyewear.

#### NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries or greenhouses.

Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment areas until spray has dried.

This product is for use on Ornamental Turf Lawns (Residential, Industrial and Institutional), Parks, Cemeteries, Athletic Fields and Golf Courses (Fairways, Aprons, Tees\* and Roughs) and similar turf areas.

\*Excluding Bentgrass Tees.

### USE PRECAUTIONS

Avoid drift of spray mist to vegetables, grapes, flowers, ornamental plants, shrubs, trees and other desirable plants. Do not pour spray solutions near desirable plants. Do not use on carpetgrass, Dichondra, not on lawns or turf where desirable clovers are present. Avoid fine mists. Use lawn type sprayer with coarse spray as wind drift is less likely. Avoid contact with exposed feeder roots of ornamentals and trees. Maximum control of weeds will be obtained from Spring or early Fall applications when weeds are actively growing. Use the higher rate for hard-to-control weeds. Do not exceed specified dosages for any area; be particularly careful within the dripline of tree and other ornamental species. Do not apply to newly seeded grasses until well established. Do not spray when air temperature exceeds 90°F.

Avoid application to Centipede or St. Augustine grasses unless turf injury can be tolerated. The suitable use of this product on non-recommended turf species may be determined by treating a small area at any rate/acre which does not exceed 3 pints/acre. Prior to treatment of larger areas, the treated areas should be observed for any sign of turf injury for a period of 30 days of normal growing conditions to determine the phytotoxicity and efficacy to the treated area. The user assumes the responsibility for any plant damage or other liability resulting from the use of this product on species not recommended on this label.

**For optimum results:** (1) Avoid applying during excessively dry or hot periods unless irrigation is used; (2) turf should be mowed 1 to 2 days before the following application; (3) reseed no sooner than 3 to 4 weeks after application of the product. Adding oil, wetting agent or other surfactant to the spray may be used to increase effectiveness on weeds, but doing so may reduce selectivity to turf resulting in turf damage. Clean and rinse spray equipment using soap or detergent and water, and rinse thoroughly before reuse for other sprays.

### WEEDS CONTROLLED BY WIL-POWER™ SELECTIVE HERBICIDE

Alder	Field pansy	Prairie
Annual yellow sweet clover	Flea bane (daisy)	Prickly lettuce
Artichoke	Flixweed	Prickly sida
Aster	Florida betony	Primrose
Austrian Fieldcress	Florida pusley	Prostrate knotweed
Bedstraw	Frenchweed	Prostrate pigweed
Beggartick	Galinsoga	Prostrate spurge
Biden	Garlic mustard	Prostrate vervain
Bindweed	Goathead	Puncture vine
Bird vetch	Goatsbeard	Purslane
Bitterweed	Goldenrod	Ragweed
Bitter wintercress	Ground ivy	Red clover
Black-eyed Susan	Gumweed	Redroot pigweed
Black medic	Hairy bittercress	Red sorrel
Black mustard	Hairy fleabane	Remstem filaree
Blackseed pantain	Hawkweed	Rough cinquefoil
Blackseed plantain	Healall	Rough fleabane
Blessed thistle	Heartleaf drymary	Roundleafed marigold
Blue lettuce	Hedge bindweed	Rush
Blue vervain	Hedge mustard	Russian pigweed
Box elder	Hemp	Russian thistle
Bracted plantain	Henbit	St. Johnswort
Brassbuttons	Hoary cress	Scarlet pimpernel
Bristly oxtongue	Hoary plantain	Scotch thistle
Broadleaf dock	Hoary vervain	Sheep sorrel
Broadleaf plantain	Honeysuckle	Shepherdspurse
Broomweed	Hop clover	Slender plantain
Buckhorn	Horsenettle	Smallflower galinsoga
Buckhorn plantain	Horsetail	Smartweed
Bulbous buttercup	Indiana mallow	Smooth dock
Bull nettle	Ironweed	Smooth pigweed
Bull thistle	Jewelweed	Sneezeweed
Burdock	Jimsonweed	Southern wild rose
Burning nettle	Kochia	Sowthistle
Bur ragweed	Knawel	Spanishneedle
Burweed	Knotweed	Spatterdock
Buttercup	Lambsquarter	Speedwell
Canada thistle	Lespedeza	Spiny
Carolina geranium	Locoweed	Spiny amaranth
Carpetweed	Lupine	Spiny cocklebur
Catchweed bedstraw	Mallow	Spotted catsear
Catsear	Marshelder	Spotted knapweed
Catnip	Matchweed	Spotted spurge
Chickweed	Mexicanweed	Spurge
Chicory	Milk vetch	Spurweed
Cinquefoil	Milkweed bloodflower	Stinging nettle
Clover	Mugwort	Stinkweed
Cockle	Morningglory	Stitchwort
Cocklebur	Mouseear chickweed	Strawberry clover
Coffebean	Musk thistle	Sumac
Coffeeweed	Mustard	Sunflower
Common chickweed	Narrowleaf plantain	Sweet clover
Common mullein	Narrowleaf vetch	Tall nettle
Common sowthistle	Nettle	Tall vervain
Corn chamomile	Orange hawkweed	Tansy mustard
Creeping jenny	Oxalis	Tansy ragwort
Crimson clover	Oxeye daisy	Tanweed
Croton	Parsley-piert	Tarweed
Cudweed	Parsnip	Thistle
Curly dock	Pearlwort	Tick trefoil
Curly indigo	Pennycress	Toadflax
Dandelion	Pennywort	Trailing crownvetch
Dead nettle	Peppergrass	Tumble mustard
Dock	Pepperweed	Tumble pigweed
Dogbane	Pigweed	Tumbleweed
Dogfennel	Pineywoods bedstraw	Velvet leaf
Elderberry	Plains coreopsis	Venice mallow
English daisy	Planatain	Veronica
Fall dandelion	Poison hemlock	Vervain
False dandelion	Poison ivy	Vetch
False flax	Poison oak	Virginia buttonweed
False sunflower	Pokeweed	Virginia creeper
Fiddleneck	Poorjoe	Virginia pepperweed
Field bindweed	Povertyweed	Wavyleaf bullthistle

**WEEDS CONTROLLED BY WIL-POWER™**

**SELECTIVE HERBICIDE Continued:**

Western clematis	Wild lettuce	Witchweed
Western salsify	Wild marigold	Woolly morningglory
White clover	Wild onion	Woodsorrel
White mustard	Wild parsnip	Woolly croton
Wild mustard	Wild radish	Woolly plantain
Wild aster	Wild rape	Wormseed
Wild buckwheat	Wild strawberry	Yarrow
Wild carrot	Wild sweet potato	Yellow rocket
Wild four-o'clock	Wild vetch	Yellowflower pepperweed
Wild garlic	Willow	and other broadleaf weeds
Wild geranium		

**ORNAMENTAL LAWNS AND TURF**

Apply Wil-Power™ at the rate of 2 to 3 pints in 20 to 240 gallons of water per acre (0.73 to 1.10 fluid ounces in 0.5 to 5.5 gallons of water per 1000 square feet) to control weeds growing in turf grasses planted to Bluegrass, Fescue, Rye, Bent (excluding golf course greens and tees), Bahia, Bermuda and Zoysia.

**Controlled Droplet Applicators – (CDA), Atomizers and Spinning Disk Applicators:** Use Wil-Power™ at the rate of 2 to 3 pints per acre (0.73 to 1.10 fluid ounces per 1000 square feet) in sufficient water to assure coverage (1 to 4 gallons of water per acre is normal for this type of equipment).

**NOTE:** For all grasses (1) Do not overlap spray patterns; (2) Use reduced rates if grass is stressed from heat, drought, etc.; and (3) Follow CDA equipment spray instructions.

**SOD FARMS**

Wil-Power™ is intended for use on sod farms. This herbicide provides selective broadleaf control in warm season and cool season turfgrass established for commercial sod production.

**Timing**

Apply this herbicide to broadleaf weeds that are actively growing. Follow-up applications may be required for dense infestations of perennial and biennial weeds. Do not apply Wil-Power™ to Floratam St. Augustinegrass.

The application of this herbicide to grass seedlings is recommended after the second mowing.

The application of Wil-Power™ to newly sodded, sprigged or plugged grasses should be delayed until 3 or 4 weeks after sodding, sprigging or plugging operations. Also, applications to dormant Bermudagrass, dormant Zoysiagrass and dormant Bahiagrass are suggested.

**Recommended Rates for Sod Farms**

For Bluegrass, Ryegrass and Fescue, apply 1-1/2 to 3 pints per acre. For Creeping Bentgrass use .9 pints per acre. For Common and Hybrid Bermuda, Bahia, Zoysia and St. Augustine grass apply 1 to 2 pints per acre. Spray volume for Sod Farm application is 5 to 175 gallons per acre.

Generally, the lower application rates will provide satisfactory control of sensitive weeds. The higher application rates will be required for dense infestations of perennial weeds, for adverse/extreme environmental conditions, or for weeds beyond the appropriate growth stages.

**COMPATIBILITY**

A compatibility test is recommended prior to mixing with liquid fertilizers and liquid iron. All regulations either state or Federal, relating to the application of liquid fertilizers or liquid iron and Wil-Power™ must be strictly followed.

**STORAGE AND DISPOSAL**

Do not contaminate water, food or feed by storage or disposal.

**PESTICIDE STORAGE:** Always use original container to store pesticides in a secured warehouse or storage building. Store at temperatures above 32° F. If allowed to freeze, remix before using. This does not alter this product. Containers should be opened in well ventilated areas. Keep container tightly sealed when not in use. Do not stack cardboard cases more than two pallets high. Do not store near open containers of fertilizer, seed or other pesticides.

**PESTICIDE DISPOSAL:** Pesticide wastes are acutely toxic. If container is damaged or if pesticide has leaked, contain all spillage. Absorb and clean up all spilled material with granules or sand. Place in a closed labeled container for proper disposal. Improper disposal of excess pesticide, spray mixtures, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

**CONTAINER DISPOSAL:** Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities. Plastic containers are also disposable by incineration, or if allowed by State and local authorities, by burning. If burned, stay out of smoke.

**CONTAINER DISPOSAL FOR REFILLABLE CONTAINERS:** Close all openings and replace all caps. Contact Nufarm Customer Service at 1-800-345-3330, to arrange for return of the empty refillable container.

**Conditions of Sale and Limitation of Warranty and Liability:**

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using the product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of the product should be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of many different factors including, without limitation, manner of use or application, weather, combination with other products, or crop conditions. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold Manufacturer and Seller harmless from all claims relating to such factors.

**Seller warrants that this product conforms to the chemical description on the label. EXCEPT FOR THIS WARRANTY, THE PRODUCT IS FURNISHED "AS-IS," AND NEITHER SELLER NOR MANUFACTURER MAKES ANY OTHER WARRANTIES, EXPRESS OR IMPLIED, WITH RESPECT TO THE SELECTION, PURCHASE OR USE OF THIS PRODUCT; SELLER AND MANUFACTURER SPECIFICALLY DISCLAIM ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE. Buyer and User accept all risks arising from any use of this product, including without limitation uses contrary to label instructions, under abnormal conditions, or under conditions not reasonably foreseeable to (or beyond the control of) Seller or Manufacturer.**

To the extent permitted by law, neither Manufacturer nor Seller shall be liable for any incidental, consequential or special damages resulting from the use or handling of this product. THE EXCLUSIVE REMEDY OF THE BUYER OR USER, AND THE EXCLUSIVE LIABILITY OF MANUFACTURER AND SELLER, FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THIS PRODUCT, OR, AT THE ELECTION OF MANUFACTURER OR SELLER, THE REPLACEMENT OF THE PRODUCT.

These Conditions of Sale and Limitation of Warranty and Liability shall be interpreted in accordance with the laws of the State of California, excluding its conflicts of laws rules, and may not be amended by any oral or written agreement.

WILBUR-ELLIS Logo®, and IDEAS TO GROW WITH® are Registered Trademarks of Wilbur-Ellis Company.

Wil-Power™ is a Trademark Licensed to Wilbur-Ellis Company.

The Optical Advantage® is a Registered Trademark of Nufarm Americas, Inc.

F-122304  
RV120104

Distributed by:  
**WILBUR-ELLIS COMPANY**  
PO BOX 16458 • FRESNO CA 93755