

Supplemental Table 2. Geometric mean and asymmetric standard error of mean (SEM) for proinsulin, 17-beta estradiol, and triglyceride concentrations

Variable	Condition	Time	Geometric mean	Positive SEM	Negative SEM
Proinsulin (pmol/L)	Placebo	0	4.8	5.4	4.3
		60	6.0	7.4	4.9
		90	21.7	27.8	17.0
		120	23.1	29.9	17.8
		150	26.9	31.0	23.2
		180	21.6	24.3	19.3
	BPA 4	0	6.2	7.2	5.3
		60	7.8	9.7	6.3
		90	18.1	24.8	13.2
		120	18.5	25.5	13.4
		150	20.2	27.9	14.6
		180	14.8	22.1	10.0
	BPA 50	0	6.2	7.6	5.1
		60	8.1	10.1	6.5
		90	28.7	32.0	25.8
		120	19.1	25.6	14.2
		150	25.3	29.9	21.3
		180	19.0	24.5	14.8
17-beta estradiol (pg/mL)	Placebo	0	51.9	65.4	41.2
		15	54.8	68.8	43.7
		30	53.6	67.5	42.7
		45	51.9	65.4	41.1
		60	54.0	69.3	42.2
		90	50.1	64.4	39.0
		120	47.5	61.5	36.7
		150	49.8	62.6	39.5
		180	51.4	63.9	41.3
	BPA 4	0	57.6	70.8	46.9
		15	57.2	71.2	45.9
		30	53.7	65.7	43.9
		45	57.4	69.8	47.1
		60	56.4	68.6	46.4
		90	52.2	63.5	42.9
		120	52.4	63.7	43.0
		150	54.1	65.8	44.5
		180	55.5	66.6	46.3
	BPA 50	0	49.9	60.9	40.8
		15	52.4	62.6	43.9
		30	50.3	60.5	41.9
		45	53.2	63.5	44.5
		60	54.0	64.9	45.0
		90	53.8	64.1	45.1
120		50.3	59.4	42.6	
150		51.6	61.6	43.3	
180		53.6	64.5	44.6	

Triglyceride (mmol/L)	Placebo	0	1.0	1.1	0.9
		15	1.0	1.1	0.9
		30	1.0	1.1	0.9
		45	1.0	1.1	0.9
		60	1.0	1.1	0.9
		90	1.1	1.1	1.0
		120	1.0	1.1	0.9
		150	0.9	1.0	0.9
		180	0.9	0.9	0.8
		BPA 4	0	1.0	1.1
	15		1.0	1.1	0.9
	30		1.0	1.1	0.9
	45		0.9	1.1	0.8
	60		0.9	1.0	0.8
	90		0.9	1.0	0.9
	120		0.9	1.0	0.8
	150		0.9	1.0	0.8
	180		0.8	0.9	0.7
	BPA 50		0	0.9	1.1
		15	1.0	1.1	0.9
30		0.9	1.1	0.8	
45		1.0	1.1	0.9	
60		1.0	1.1	0.8	
90		1.0	1.1	0.9	
120		0.9	1.1	0.8	
150		0.9	1.0	0.8	
180		0.8	0.9	0.7	

---